

NPS-OR-93-017

NAVAL POSTGRADUATE SCHOOL

Monterey, California





ASYMPTOTIC PROPERTIES OF STOCHASTIC GREEDY BIN-PACKING

93-29562

Donald P. Gaver Patricia A. Jacobs

November 1993

Approved for public release; distribution is unlimited.

Prepared for: Naval Postgraduate School Monterey, CA 93943

NAVAL POSTGRADUATE SCHOOL MONTEREY, CA 93943-5000

Rear Admiral T. A. Mercer Superintendent

Harrison Shull Provost

This report was prepared in conjunction with research funded by the Naval Postgraduate School Direct Funded Research Program.

Reproduction of all or part of this report is authorized.

This report was prepared by:

DONALD P. GAVER, JR.

Professor of Operations Research

PATRICIA A. JACOBS

Professor of Operations Research

Reviewed by:

PETER PERDITE

Professor and Chairman

Department of Operations Research

Released by:

PAUL J. MÄRTQ

Dean of Research

Public reporting burden for this collection of a gathering and mentairing the data needed, collection of information, including suggestion Davis Highway, Suite 1204 Artinoton, VA	information is estimated to average 1 hour per and completing and reviewing the collection in ns for reducing this burden, to Washington He 22202-4302, and to the Office of Manageme	response, including the time for rewood information. Send comments regal acquarters Services. Directorate for int and Budget, Paperwork Reduction	wing instructions, searching existing data sources, rding this burden estimate or any other aspect of this information Operations and Reports, 1215 Jefferson in Project (0704-0188), Washington, DC 20503
1. AGENCY USE ONLY (Leave blank)		3. REPORT TYPE AND DAT	
The second second second second	November 1993	Technical Rep	
4. TITLE AND SUBTITLE		- 	S. FUNDING NUMBERS
	of Stochastic Greedy Bi	n-Packing	ORGV1
6. AUTHOR(S)		1	3153
Donald P. Gaver and I	Patricia A. Jacobs		
7. PERFORMING ORGANIZATION NAM	ME(S) AND ADDRESS(ES)		B. PERFORMING ORGANIZATION REPORT NUMBER
Navel Postere dunta Co	ah a al		NEI ON MONDEY
Naval Postgraduate So	cnool	J	NPS-OR-93-017
Monterey, CA 93943			
9. SPONSORING / MONITORING AGE	NCY NAME(S) AND ADDRESS(ES)		10. SPONSORING / MONITORING AGENCY REPORT NUMBER
Naval Postgraduate So	chool	1	
Monterey, CA 93943		ļ	
11. SUPPLEMENTARY NOTES			
128. DISTRIBUTION / AVAILABILITY	STATEMENT		12b. DISTRIBUTION CODE
Approved for public r	elease, distribution is un	limited.	
13. ABSTRACT (Maximum 200 wor	ds)		
attention of a number, contents become nearl jobs becomes large. Exasymptotics are compared to the compared to	I, of processors is studie y jointly but degenerated plicit and simple paramared with simulation. The cess is quantified, and se	ed. It is shown that the control of	r scheduling jobs for the he suitably normalized bin if the rate of approach of as are supplied and the greedy policy over a laissez-
14. SUBJECT TERMS			15. NUMBER OF PAGES
greedy algorithm, bin	27		
Ornstein-Uhlenbeck p		, <u>,</u>	16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	ON 20. LIMITATION OF ABSTRACT
Unclassified	Unclassified	Unclassified	UL

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

ASYMPTOTIC PROPERTIES OF STOCHASTIC GREEDY BIN-PACKING

D. P. Gaver P. A. Jacobs

DITIC QUALITY HISPEGUED 5

Accesion For NTIS OPERAL DTIC CALL U. and Call Call Ey Color Call Available Call Special A-1

Abstract

An adaptive or greedy policy for packing I bins, or equivalently for scheduling jobs for the attention of a number, I, of processors is studied. It is shown that the suitably normalized bin contents become nearly jointly but degenerately Gaussian/normal if the rate of approach of jobs becomes large. Explicit and simple parameter characterizations are supplied and the asymptotics are compared with simulation. The advantage of the greedy policy over a laissez-faire policy of equal access is quantified, and seen to depend upon $\sqrt{\text{number of bins or processors}}$.

Introduction

We study an adaptive or greedy policy for packing *I* bins, or equivalently for scheduling jobs for the attention of a number, *I*, of processors. The connection between bin-packing and makespan scheduling is well described in Coffman and Lueker [1], chapter 1.

It is shown that the suitably normalized bin contents become nearly jointly but degenerately Gaussian/normal if the rate of approach of jobs becomes large. Explicit and simple parameter characterizations are supplied, and the asymptotics are compared with simulation. The advantage of the greedy policy over a laissez-faire policy of equal access is quantified, and seen to depend upon $\sqrt{\text{number of bins or processors}}$.

Greedy Bin-Filling

There are I bins (potential servers). Jobs arrive at the bin system according to a homogeneous Poisson process, rate λ . Each job size is independently and identically distributed according to $F_B(x)$; a generic job size is B, a real positive random variable. Let $N_i(t)$ denote amount of work in bin i at time t; this is the sum of the job sizes deposited in that bin up to time t. We refer to $N_i(t)$ as the bin size at time t hereafter, although other terminology is used in the scheduling literature.

Consider *Policy* E (Equalization): When a new job arrives it is deposited in the bin with smallest amount of accumulated work. $N(t) = \{N_i(t), t \ge 0, i \in (1, 2, ..., I)\}$ is a Markov process with the following generator; for $\Delta > 0$

$$N_i(t + \Delta) = N_i(t)$$
 with probability $1 - \lambda \Delta p_i(N_i(t), N(t)) + o(\Delta)$
$$= N_i(t) + B(t) \text{ with probability } \lambda \Delta p_i(N_i(t), N(t)) + o(\Delta)$$
 (1,

Define

$$p_i(N_i(t), N(t)) = \begin{cases} 1 & \text{if } N_i(t) \le N_j(t), j \ne i \\ 0 & \text{otherwise} \end{cases}$$

Consider also Probabilistic Policy E: same as above but

$$p_i(N_i(t), N(t)) = \frac{h(N_i(t))}{\sum_{j=1}^{i} h(N_j(t))}$$
(2)

where $h(\cdot)$ is a positive homogeneous function chosen to become large when $N_i(t)$ is small; e.g. $h(x) = x^{-p}$. Such a policy leads to workload growth very similar to Policy E's, but can be analyzed more readily. See Gaver, Morrison and Silveira [2] for application of such a probabilistic policy in a service-system scheduling context. In the present context, the bins can be buffers containing jobs to be

processed later and the aim is to keep the total processing time short; it is (optimistically) assumed that the contents of each bin is known at all times to the scheduler and that each processing time is known when the job or task appears.

Joint Moment-Generating Function

Let the moment-generating function (assumed to exist, otherwise use the characteristic function) be

$$\psi(\theta,t) \stackrel{(d)}{=} E\left[e^{\theta \cdot N(t)}\right] = E\left[\exp\left(\sum_{j=1}^{l} \theta_{j} N_{j}(t)\right)\right]. \tag{3}$$

Condition on $N_i(t)$, $i \in (1, 2, ..., I)$ and use the generator to obtain

$$\begin{split} E &\left[\exp \left(\sum_{j=1}^{I} \theta_{j} N_{j}(t + \Delta) \right) \middle| N(t), B(t) \right] \\ &= \left[1 - \lambda \Delta \right] \exp \left(\sum_{j=1}^{I} \theta_{j} N_{j}(t) \right) + \\ &+ \lambda \Delta \sum_{j=1}^{I} \left[\exp \left(\theta_{j} \left(N_{j}(t) + B(t) \right) + \sum_{k \neq j} \theta_{k} N_{k}(t) \right) \middle| p_{j} \left(N_{j}(t); N(t) \right) \right] + o(\Delta). \end{split}$$

Remove conditions, defining the m g f of the task size arriving at t, B(t), to be $\hat{b}(\theta)$, to find

$$\psi(\theta, t + \Delta) = (1 - \lambda \Delta)\psi(\theta, t) + \lambda \Delta \sum_{j=1}^{J} \hat{b}(\theta_j) E\left[\exp(\theta_j N_j(t)) \cdot p_j(N_j(t); N(t))\right] + o(\Delta).$$

Let $\Delta \rightarrow 0$ to get

$$\frac{\partial \psi(\theta,t)}{\partial t} = -\lambda \psi(\theta,t) + \lambda \sum_{j=1}^{I} \hat{b}(\theta_j) E\left[\exp(\theta_j N_j(t)) \cdot p_j(N_j(t); N(t))\right]. \tag{4}$$

Note that nothing that follows prevents non-stationary input rates: i.g. $\lambda = \lambda r(t)$, and the mgf of B(t) to be $\hat{b}(\theta;t)$ provided these do not drop quickly to zero.

Scaling

Put

$$X_{j}(t) = \frac{N_{j}(t) - \lambda \beta_{j}(t)}{\sqrt{\lambda}}$$
 (5)

and let $\lambda \gg 1$. It is anticipated that with suitable choice of the functions $\{\beta_j(t)\}$ $\{X_j(t), j \in (1, 2, ..., I)\}$ should become a Gaussian process as $\lambda \to \infty$. Let $\varphi(\theta, t; \lambda) = E\left[\exp\left(\sum_{j=1}^{I} \theta_j X_j(t)\right)\right]$ for $X_j(t)$ defined as above. Note that

$$E\left[\exp\left(\sum_{j=1}^{I} (\theta_{j} / \sqrt{\lambda}) N_{j}(t)\right)\right] = \psi(\theta / \sqrt{\lambda}, t)$$

$$= E\left[\exp\left(\sum_{j=1}^{I} \theta_{j} X_{j}(t)\right)\right] \cdot \exp\left(\sqrt{\lambda} \sum_{j=1}^{I} \theta_{j} \beta_{j}(t)\right)$$

$$= \varphi(\theta, t; \lambda) \exp(\sqrt{\lambda} \theta \beta(t)).$$
(6)

Now since from (4)

$$\frac{\partial \psi(\theta / \sqrt{\lambda}, t)}{\partial t} = -\lambda \psi(\theta / \sqrt{\lambda}, t) + \lambda \sum_{j=1}^{I} \hat{b}(\theta_j / \sqrt{\lambda}) E\left[e^{\theta_j / \sqrt{\lambda} N_j(t)} \cdot p_j(N_j(t); N(t))\right],$$

substitution of (5) yields

$$\frac{\partial \psi(\theta / \sqrt{\lambda}, t)}{\partial t} = \frac{\partial}{\partial t} \Big[\psi(\theta, t; \lambda) \exp(\sqrt{\lambda} \theta \beta(t)) \Big] = \frac{\partial \psi(\theta, t; \lambda)}{\partial t} \exp(\sqrt{\lambda} \theta \beta(t)) \\
+ \psi(\theta, t; \lambda) \Big(\sqrt{\lambda} \theta \beta'(t) \exp(\sqrt{\lambda} \theta \beta(t)) \Big) \\
= -\lambda \psi(\theta, t; \lambda) \exp(\sqrt{\lambda} \theta \beta(t)) + \\
+ \lambda \sum_{j=1}^{I} \hat{b} \Big(\theta_j / \sqrt{\lambda} \Big) E \Big[\exp(\theta X(t)) \cdot \exp(\sqrt{\lambda} \theta \beta(t)) \cdot p_j \Big(\lambda \beta_j(t) + \sqrt{\lambda} X_j(t); \lambda \beta + \sqrt{\lambda} X \Big) \Big].$$

Cancel $\exp(\sqrt{\lambda}\theta\beta(t))$ to obtain

$$\frac{\partial \varphi(\theta, t; \lambda)}{\partial t} + \varphi(\theta, t; \lambda) \sqrt{\lambda} \theta \beta'(t) = -\lambda \varphi(\theta, t; \lambda)$$

$$+ \lambda \sum_{j=1}^{l} \hat{b} \Big(\theta_j / \sqrt{\lambda} \Big) E \Big[\exp(\theta X(t)) \cdot p_j \Big(\lambda \beta_j(t) + \sqrt{\lambda} X_j(t); \lambda \beta + \sqrt{\lambda} X \Big) \Big]. \tag{7}$$

Let

$$p_{j}(N_{j}(t),N(t)) = \frac{h(N_{j}(t))}{\sum_{k=1}^{I} h_{k}(N_{k}(t))} = \frac{h(\lambda \beta_{j}(t) + \sqrt{\lambda} X_{j}(t))}{\sum_{k=1}^{I} h(\lambda \beta_{k}(t) + \sqrt{\lambda} X_{k}(t))}$$

where $h(\cdot)$ is homogeneous: $h(\lambda x) = \lambda Ph(x)$. Consequently

$$p_{j}\left(\lambda\beta_{j}(t) + \sqrt{\lambda}X_{j}(t); \lambda\beta(t) + \sqrt{\lambda}X(t)\right) = \frac{h\left(\beta_{j}(t) + X_{j}(t) / \sqrt{\lambda}\right)}{\sum_{k=1}^{J} h\left(\beta_{k}(t) + X_{k}(t) / \sqrt{\lambda}\right)}$$
(8)

(h(x)) assumed to be differentiable). Expand in inverse powers of $\sqrt{\lambda}$:

$$p_j\Big(\lambda\beta_j(t)+\sqrt{\lambda}X_j(t);\lambda\beta(t)+\sqrt{\lambda}X(t)\Big)=$$

$$\frac{h(\beta_{j}(t))}{\sum\limits_{k=1}^{I}h(\beta_{k}(t))} + \frac{1}{\sqrt{\lambda}} \left[\frac{h'(\beta_{j}(t))X_{j}(t)}{\sum\limits_{k=1}^{I}h(\beta_{k}(t))} - \frac{h(\beta_{j}(t))}{\left(\sum\limits_{k=1}^{I}h(\beta_{k}(t))\right)^{2}} \sum_{k=1}^{I}h'(\beta_{k}(t))X_{k}(t) \right] + O\left(\frac{1}{\lambda}\right). \tag{9}$$

Since

$$\sum_{j=1}^{I} p_{j} \left(\lambda \beta_{j}(t) + \sqrt{\lambda} X_{j}(t); \lambda \beta(t) + \sqrt{\lambda} X(t) \right) = 1$$

$$= \sum_{j=1}^{I} \frac{h(\beta_{j}(t))}{\sum_{k=1}^{I} h(\beta_{k}(t))}$$
(9,a)

this implies that the summed coefficients of $1/\sqrt{\lambda}$, $1/\lambda$, etc. must be individually zero; this is easily verified for the coefficient of $1/\sqrt{\lambda}$.

Asymptotic Expansion For $\varphi(\theta,t;\lambda)$.

Put

$$\varphi(\theta,t;\lambda) = \varphi_0(\theta,t) + \frac{1}{\sqrt{\lambda}} \varphi_1(\theta,t) + \frac{1}{\lambda} \varphi_2(\theta,t) + \dots$$

This can now be entered into (7) and evaluated by means of (9):

$$\sum_{\ell=0}^{\infty} \frac{\partial \varphi_{\ell}}{\partial t} \cdot \frac{1}{\left(\sqrt{\lambda}\right)^{\ell}} + \sqrt{\lambda} \,\theta \beta'(t) \sum_{\ell=0}^{\infty} \varphi_{\ell} \cdot \frac{1}{\left(\sqrt{\lambda}\right)^{\ell}} = -\lambda \sum_{\ell=0}^{\infty} \varphi_{\ell} \frac{1}{\left(\sqrt{\lambda}\right)^{\ell}}$$

$$+\lambda \sum_{j=1}^{I} \left(\sum_{k=0}^{\infty} b_{k} \frac{1}{k!} \,\theta_{j}^{k} / \left(\sqrt{\lambda}\right)^{k}\right) \left[\left(\sum_{\ell=0}^{\infty} \varphi_{\ell} \frac{1}{\left(\sqrt{\lambda}\right)^{\ell}}\right) \left(\frac{h(\beta_{j}(t))}{\sum_{k=1}^{I} h(\beta_{k}(t))}\right)\right]$$

$$(10)$$

$$+\frac{1}{\sqrt{\lambda}}\left(\frac{h'(\beta_{j}(t))}{\sum_{k=1}^{I}h(\beta_{k}(t))}\left(\sum_{\ell=0}^{\infty}\frac{\partial\varphi_{\ell}}{\partial\theta_{j}}\frac{1}{\left(\sqrt{\lambda}\right)^{\ell}}\right)-\frac{h(\beta_{j}(t))}{\left(\sum_{k=1}^{I}h(\beta_{k}(t))\right)^{2}}\sum_{k=1}^{I}h'(\beta_{k}(t))\sum_{\ell=0}^{\infty}\frac{\partial\varphi_{\ell}}{\partial\theta_{k}}\frac{1}{\left(\sqrt{\lambda}\right)^{\ell}}\right)+O\left(\frac{1}{\lambda}\right)\right]$$

In the above $b_k = E[B^k]$, the k^{th} moment of job size; of course $b_0 = 1$.

Now identify the coefficients of inverse powers of $\sqrt{\lambda}$, and thereby equations for φ_{ℓ} and β_{ℓ} . From (10) for $\ell=0$,

$$\frac{\partial \varphi_0}{\partial t} + \sqrt{\lambda} \theta \beta'(t) \cdot \varphi_0 = -\lambda \varphi_0 + \lambda \sum_{j=1}^{I} \left(1 + b_1 \theta_j \frac{1}{\sqrt{\lambda}} + \frac{1}{2} b_2 \theta_j^2 \frac{1}{\lambda} + \dots \right)$$

$$\left[\varphi_{0} \cdot \frac{h(\beta_{j}(t))}{\sum_{k=1}^{I} h(\beta_{k}(t))} + \frac{1}{\sqrt{\lambda}} \left(\frac{h'(\beta_{j}(t))}{\sum_{k=1}^{I} h(\beta_{k}(t))} \frac{\partial \varphi_{0}}{\partial \theta_{j}} - \frac{h(\beta_{j}(t))}{\left[\sum_{k=1}^{I} h(\beta_{k}(t))\right]^{2}} \sum_{k=1}^{I} h'(\beta_{k}(t)) \frac{\partial \varphi_{0}}{\partial \theta_{k}} \right] + O\left(\frac{1}{\lambda}\right) \right]$$
(11)

The terms of order λ cancel from the r h s. The terms of order $\sqrt{\lambda}$ on l h s and r h s cancel if

$$\theta \beta'(t) \varphi_0 = b_1 \sum_{j=1}^{l} \theta_j \frac{h(\beta_j(t))}{\sum_k h(\beta_k(t))} \cdot \varphi_0.$$

In order for this to occur,

$$\frac{d\beta_j}{dt} = b_1 \frac{h(\beta_j(t))}{\sum_k h(\beta_k(t))}, \quad j = 1, 2, \dots, l$$
 (12)

the solution of which determines $\beta_i(t)$.

Next look for terms of order 1. The l h s provides $\frac{\partial \varphi_0}{\partial t}$. The r h s provides the

terms
$$\varphi_0 \sum_{j=1}^{I} b_2 \frac{\theta_j^2}{2} \frac{h(\beta_j(t))}{\sum_k h(\beta_k(t))}$$

and
$$b_1 \sum_{j=1}^{l} \theta_j \left(\frac{h'(\beta_j(t))}{\sum_{k} h(\beta_k(t))} \frac{\partial \varphi_0}{\partial \theta_j} - \frac{h(\beta_j(t))}{\left(\sum_{k} h(\beta_k(t))\right)^2} \sum_{k=1}^{l} h'(\beta_k(t)) \frac{\partial \varphi_0}{\partial \theta_k} \right)$$

Note that the condition (12) actually annihilates any term of order 1 (or higher) in φ_{ℓ} for $\ell = 1, 2, ...$ on the r h s, and the discussion of (9, a) shows that there is no contribution from $\partial \varphi_{\ell}/\partial \theta_{j}$. Consequently

$$\frac{\partial \varphi_{0}}{\partial t} = \frac{b_{2}}{2} \sum_{j=1}^{I} \theta_{j}^{2} p_{j} (\beta_{j}(t)) \varphi_{0}(\theta, t) +$$

$$b_{1} \left[\sum_{j=1}^{I} \theta_{j} \left(\frac{h'(\beta_{j}(t))}{\sum_{k} h(\beta_{k}(t))} \frac{\partial \varphi_{0}(\theta, t)}{\partial \theta_{j}} - \frac{h(\beta_{j}(t))}{\sum_{k} h(\beta_{k}(t))} \right)^{2} \sum_{k}^{I} h'(\beta_{k}(t)) \frac{\partial \varphi_{0}(\theta, t)}{\partial \theta_{k}} \right]$$

$$\equiv \frac{b_{2}}{2} \sum_{j=1}^{I} \theta_{j}^{2} p_{j} (\beta_{j}(t)) \varphi_{0}(\theta, t) + b_{1} \sum_{j=1}^{I} \theta_{j} \left(H_{j}(t) \frac{\partial \varphi_{0}}{\partial \theta_{j}} - \sum_{k} H_{jk}(t) \frac{\partial \varphi_{0}(\theta, t)}{\partial \theta_{k}} \right)$$

$$\text{Note: } \sum_{j=1}^{I} \left(H_{j}(t) \frac{\partial \varphi_{\ell}}{\partial \theta_{j}} \frac{1}{(\sqrt{\lambda})^{\ell}} - \sum_{k} H_{jk}(t) \frac{\partial \varphi_{\ell}}{\partial \theta_{k}} \frac{1}{(\sqrt{\lambda})^{k}} \right) = 0.$$

The PDE for $\varphi_0(\theta,t)$ is recognizable as that of an Ornstein-Uhlenbeck (Gaussian) process. Similar equations for higher-order corrections can be derived similarly, but we omit this step.

MOMENTS

The (0^{th} order) joint moments of $X(t) = (X_j(t), j = 1, 2, ..., I)$ satisfy ordinary differential equations that are readily obtained by differentiation of (13) at $\theta = 0$. If $V_i(t) = E[X_i^2(t)]$, $V_{ij}(t) = E[X_i(t)X_j(t)]$ for $i \neq j$ then

$$\frac{dV_{i}}{dt} = b_{2}p_{i}(\beta_{i}(t)) + 2b_{1} \left[H_{i}(t)V_{i}(t) - \sum_{(k)} H_{ik}(t)V_{ik}(t) \right]
\frac{dV_{ij}}{dt} = b_{1} \left[\left(H_{i}(t) + H_{j}(t) \right) V_{ij}(t) - \left(\sum_{(k)} H_{jk}(t)V_{ik}(t) + \sum_{(k)} H_{ik}(t)V_{jk}(t) \right) \right]$$
($i \neq i$)

The above must be tailored as follows:

$$\frac{dV_{i}(t)}{dt} = b_{2}p_{i}(\beta_{i}(t)) + 2b_{1}\left[(H_{i}(t) - H_{ii}(t))V_{i}(t) - \sum_{k \neq i} H_{ik}(t)V_{ik}(t) \right]$$
(14)

$$\frac{dV_{ij}}{dt} = b_1 \begin{bmatrix} \left(H_i(t) + H_j(t) \right) V_{ij}(t) - \left(H_{ji} V_i(t) + H_{ij} V_j(t) \right) \\ - \left(\sum_{(k \neq i)} H_{jk}(t) V_{ik}(t) + \sum_{(k \neq j)} H_{ik}(t) V_{jk}(t) \right) \end{bmatrix}$$

Now return to specifics; consider (12): for $j \neq k$

$$\frac{d\beta_{j}(t)}{h(\beta_{j}(t))} = \frac{d\beta_{k}(t)}{h(\beta_{k}(t))}$$

if the bins are filled as suggested. This implies that

$$\beta_{j}(t) = \frac{b_{1}t}{I}, \text{ all } j \in (1, 2, ..., I)$$
 (15)

and $p_I(\beta_I(t)) = 1/I$.

For the case in which $h(x) = x^{-p}$ it can be seen that

$$H_i(t) = -\frac{p}{b_1 t}, \quad H_{ij}(t) = -\frac{p}{Ib_1 t}$$
 (16)

Substitute into (14) to obtain these equations for $V_i(t) \equiv V(t)$, $(\forall i)$;

 $V_{ij}(t) = W(t), (\forall i \neq j)$:

$$\frac{dV}{dt} = \frac{b_2}{I} - \frac{2p}{t} \left(\frac{I-1}{I}\right) (V-W) \tag{17,a}$$

$$\frac{dW}{dt} = \frac{2p}{lt}(V - W),\tag{17,b}$$

from which an equation for Z(t) = V(t) - W(t) emerges:

$$\frac{dZ}{dt} + \frac{2p}{t}Z(t) = \frac{b_2}{I} \tag{18}$$

A solution to (18) over $t \in (L, \infty)$, L > 0 is of this form:

$$Z(t) = \frac{b_2t}{I(1+2p)} + \frac{K(L)}{t^{2p}}, \qquad L \le t$$

From this and (17,b)

$$\frac{dW(t)}{dt} = \frac{2p}{lt}Z(t)$$

we get

$$W(t) = \frac{2p}{1+2p} \frac{b_2}{I^2} t + \frac{K(L)}{2p} \left(\frac{1}{t^{2p}} - \frac{1}{L^{2p}} \right) \qquad L \le t$$
 (19)

and from (17,a)

$$\frac{dV(t)}{dt} = \frac{b_2}{I} - \frac{2p}{t} \left(\frac{I-1}{I}\right) Z(t)$$

$$V(t) = \frac{b_2 t}{I^2} \left(\frac{2p+I}{2p+1} \right) + K(L) \left(\frac{I-1}{I} \right) \left(\frac{1}{t^{2p+1}} - \frac{1}{L^{2p+1}} \right)$$
 (20)

Now if p >> 1 it is seen that

$$V(t) = W(t) = \frac{b_2 t}{I^2} \tag{21}$$

Parenthetically, the comparable figures for independently filled bins are

$$V(t) = \frac{b_2 t}{I}$$
 and $W(t) = 0$. (22)

From (15) and (20)

$$E[N_i(t)] = \lambda t \frac{b_1}{I} + O(\sqrt{\lambda})$$

$$Var[N_i(t)] = Cov[N_i(t), N_j(t)] = \lambda t \frac{b_2}{l^2} + O(\sqrt{\lambda})$$
(23)

The singular behavior of W(t) and V(t) for small t, as in (19) and (20), can be attributed to the indeterminacy of the bin selection probability, (2), for $N_i(0) = 0$, \forall_i , which was the assumed initial condition. The long-time behavior of greedy packing, expressed by (23), is of interest: since $Var[N_i(t)] = Cov[N_i(t), N_j(t)]$ all bin contents are essentially perfectly correlated at any time. Consequently, to order $\sqrt{\lambda}$ the maximum bin contents $N_m(t)$, are approximately normal/Gaussian with mean $\lambda tb_1/I$ and standard deviation $\sqrt{\lambda tb_2}$ / I. If bins are filled independently the mean is the same but now $N_m(t)$ is distributed approximately as the

maximum of I independent normals, each with standard deviation $\sqrt{\lambda t b_2}$ / \sqrt{I} — considerably larger in a probabilistic sense. Note that putting p=0 in (19), (20) yields the independent result; putting p finite yields other probabilistic options. In the scheduling context the makespan, i.e. time to complete all tasks present at time t, is substantially reduced by the current greedy scheme, which is equivalent to what Coffman et al. (1991) call list scheduling (LS); our approach is on-line list scheduling, meaning that tasks are assigned to processors sequentially as they appear in time. It should be pointed out that the moments obtained above can also be derived directly from (1) and (2) by expansion of (5), rather as suggested by Isham [3].

Simulations

Limited informal simulations were conducted in order to check the accuracy of the proposed asymptotic approximations. The simulations were written in APL2 and conducted on an AMDAHL 5995–700A at the Naval Postgraduate School using the LLRANDOM random number generator; cf. Lewis et al. [4]. All simulations were run for time t=1 at the indicated λ -values for two job size distributions, both gamma: the exponential and an extended-tail highly-skewed gamma with shape parameter one-half.

Examination of Table 1 indicates that agreement is good between the asymptotic approximation and simulation results (based on 1000 replications) for the marginal distributional properties of an arbitrary bin when the greedy policy is followed. As anticipated, a considerable reduction in the variance of bin size, and also of upper-tail percentiles, is achieved by greediness, as contrasted to a simple random assignment.

The figures of Table 2, which describe the approximation to the maximum bin size, or makespan in a scheduling context, are serviceable but tend to be low or

optimistic, especially for the smaller λ -value of 50. For λ = 300 the agreement is better and correctly predicts the substantial reduction of mean, variance, and upper percentiles achieved by the greedy policy. Note that numerical agreement between simulation and our asymptotics should improve if the job sizes have smaller variances and third moments.

It can be conjectured, and demonstrated, that a cyclic or round-robin policy of putting every I^{th} arrival in the same bin will tend to reduce within-bin variance and makespan levels. It may be advantageous that both random and round-robin policies require no information concerning current bin size or occupancy at the time an assignment must be made, whereas the greedy policy and others depend on precise distributional forms do require such information. If reduction of bin size or makespan variation is important the acquisition of the information needed to implement a greedy policy may be well worth the cost.

Acknowledgments.

The writers are grateful to Paul Wright and Ed Coffman of AT&T Bell Laboratories, Murray Hill, for acquainting us with the greedy algorithm evaluation problem.

Table 1. ARBITRARY BIN SIZE

I = 5

B: G = 1: Gamma, Shape Param. = 1 (Exponential) G = 0.5: Gamma, Shape Param. = 0.5; E[B] = 1. Number Replications = 1000

					Po	licy			
			Greed	ly			Rand	dom	
1				Qua	ntiles		Quanti		
Mean Demand(λ)	Job Size (B)	Mean	Var	8 0%	90%	Mean	Var	80%	90%
50	G = 1	App: 10.0	4.0	11.7	12.6	10.0	20.0	13.8	15.7
		Sim: 10.1	5.0	11.9	13.0	10.0	19.6	13.7	16.1
	G = 0.5	App: 10.0	6.0	12.3	13.1	10.0	30.0	14.6	17.0
		Sim: 10.0	8.5	12.1	13.8	10.2	32.6	14.8	18.0
3 00	G = 1	App: 60.0	24.0	64.1	66.3	60.0	120.0	69.2	74.0
 		Sim: 60.2	24.1	64.4	66.4	60.4	112.1	69.5	73.9
	G = 0.5	App: 60.0	36.0	65.0	67.7	60.0	180.0	71.2	<i>7</i> 7.1
		Sim: 60.2	38.9	65.4	68.0	60.0	172.7	70.7	<i>7</i> 7.1

Table 2. MAXIMUM BIN SIZE

I = 5

G = 1: Gamma, Shape Param. = 1 (Exponential) G = 0.5: Gamma, Shape Param. = 0.5; E[B] = 1. Number Replications = 1000 **B**: G = 1:

					Po	licy			
1			Greed	ly			Rand	iom	l
				Qua	ntiles			Quantiles	
Mean Demand(λ)	Job Size (<i>B</i>)	Mean	Var	80%	90%	Mean	Var	80%	90%
50	G = 1	App: 10.0	4.0	11.7	12.6	—	_	17.6	19.1
		Sim: 11.2	5.2	13.1	14.2	15.6	15.1	18.7	23.9
	G = 0.5	App: 10.0	6.0	12.1	13.1	_	_	19.4	21.2
		Sim: 12.2	10.7	14.8	16.5	16.9	28.2	20.7	23.9
300	G = 1	App: 60.0	24.0	64.1	66.3	_	_	78.7	82.3
		Sim: 61.5	24.3	65.6	67.7	73.3	61.5	79.5	83.5
	G = 0.5	App: 60.0	36.0	65.0	67.7	_		82.9	87.3
		Sim: 62.3	40.0	67.7	70.3	76.2	107.6	84.3	90.4

^{—:} Not convenient to compute

REFERENCES

- [1] Coffman Jr., E. G. and Lueker, G. S. (1991). Probabilistic Analysis of Packing and Partitioning Algorithms. Wiley-Interscience. John Wiley and Sons, Inc. New York.
- [2] Gaver, D. P., Morrison, J. A., and Silveira, R. (1993). Service-adaptive multitype repairman problems. SIAM J. Appl. Math., Vol. 53, No. 2, pp. 459-470.
- [3] Isham, V. (1991). Assessing the variability of stochastic epidemics. Mathematical Biosciences. Vol. 107, pp. 209-224.
- [4] Lewis, P. A. W. and Uribe, L. (1981). "The New Naval Postgraduate School Random number generator LLRANDOMII." Naval Postgraduate School Technical Report NPS 55-81-005, Monterey, CA.

INITIAL DISTRIBUTION LIST

1.	Library (Code 052)2 Naval Postgraduate School Monterey, CA 93943-5000
2.	Defense Technical Information Center
3.	Research Office (Code 08)
4.	Prof. Peter Purdue
5.	Prof. Michael Bailey
6.	Department of Operations Research (Code OR)
7 .	Prof. Donald Gaver, Code OR-Gv
8.	Prof. Patricia Jacobs
9.	Center for Naval Analyses

10.	Dr. J. Abrahams, Code 1111, Room 6071
	Mathematical Sciences Division, Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5000
11.	Dr. David Brillinger
12.	Dr. David Burman
13.	Prof. H. Chernoff
14.	Dr. Edward G. Coffman, Jr
15.	Professor Sir David Cox
16.	Professor H. G. Daellenbach
17.	Dr. S. R. Dalal

18.	Dr. D. F. Daley
19.	Dr. B. Doshi
2 0.	Prof. Bradley Efron
21.	Dr. Guy Fayolle
22.	Dr. M. J. Fischer
23.	Prof. George S. Fishman
24.	Dr. Neil Gerr
25.	Dr. R. J. Gibbens

26.	Prof. Peter Glynn
27.	Prof. Linda V. Green
28.	Dr. Shlomo Halfin
29 .	Prof. Bernard Harris
30.	Dr. P. Heidelberger
31.	Prof. J. Michael Harrison
32.	Arthur P. Hurter, Jr
33.	Prof. D. L. Iglehart

34.	Institute for Defense Analysis
35.	Prof. J. B. Kadane
36.	Dr. F. P. Kelly
37.	Dr. Jon Kettenring
38.	Koh Peng Kong
39.	Dr. Prabha Kumar
40.	Dr. A. J. Lawrence
41.	Prof. M. Leadbetter

42.	Prof. J. Lehoczky
43 .	Dr. D. M. Lucantoni
44.	Dr. Colin Mallows
4 5.	Prof. R. Douglas Martin Department of Statistics, GN-22 University of Washington Seattle, WA 98195
46.	Prof. M. Mazumdar
47.	Dr. James McKenna
48.	Operations Research Center, Rm E40-164
49.	Dr. John Orav

50.	Dr. T. J. Ott
51.	Prof. Paul Moose
52.	Dr. John A. Morrison
53.	Dr. V. Ramaswami
54.	Dr. Martin Reiman 1 Rm #2C-117 AT&T Bell labs 600 Mountain Ave. Murray Hill, NJ 07974-2040
55.	Prof. Maria Rieders
56.	Dr. John E. Rolph
57.	Prof. M. Rosenblatt

58.	Statistics Department University of California Davis, CA 95616
59.	Prof. W. R. Schucany
60.	Dr. Rhonda Righter
61.	Prof. G. Shantikumar
62.	Prof. D. C. Siegmund 1 Dept. of Statistics Sequoia Hall Stanford University Stanford, CA 94305
63.	Prof. N. D. Singpurwalla
64.	Prof. H. Solomon
65 .	Prof. J. R. Thompson

66.	Statistics Dept., Fine Hall Princeton University Princeton, NJ 08540
67.	Dr. D. Vere-Jones
68.	Dr. Daniel H. Wagner
69 .	Dr. Ed Wegman
70.	Dr. Alan Weiss
7 1.	Dr. P. Welch
72 .	Prof. Roy Welsch